

Quellenverzeichnis

Die Bienen-Apotheke – Propolis, Honig & Co. – Das Heilwissen aus dem Bienenstock

© 2026 Markus Lischwe · Lischwe Media

Dieses Verzeichnis enthält alle 66 in den Fußnoten der gedruckten Ausgabe zitierten Quellen, nach Kapiteln gruppiert. Die Nummerierung [1]...[N] entspricht der durchlaufenden Fußnoten-Nummer im Buch. Alle verfügbaren DOIs und Weblinks sind klickbar – ein Klick führt direkt zum Volltext bzw. zur Verlagsseite.

Kapitel 1: Warum Bienen die ältesten Apotheker der Welt sind

- [1] Cremer, S. et al. (2007): Social immunity. *Current Biology*, 17(16), R693–R702.
- [2] Evans, J. D. & Spivak, M. (2010): Socialized medicine: individual and communal disease barriers in honey bees. *Journal of Invertebrate Pathology*, 103, S. S62–S72. → doi.org/10.1016/j.jip.2009.06.019
- [3] Starks, P. T. et al. (2000): Fever in honeybee colonies. *Naturwissenschaften*, 87(5), S. 229–231. → doi.org/10.1007/s001140050709
- [5] Simone-Finstrom, M. & Spivak, M. (2010): Propolis and bee health: the natural history and significance of resin use by honey bees. *Apidologie*, 41(3), S. 295–311. → doi.org/10.1051/apido/2010016
- [7] Simone-Finstrom, M. & Spivak, M. (2012): Increased resin collection after parasite challenge: a case of self-medication in honey bees? *PLOS ONE*, 7(3), e34601. → doi.org/10.1371/journal.pone.0034601
- [8] Borba, R. S. et al. (2015): Propolis envelope in *Apis mellifera* colonies supports honey bees against the pathogen *Paenibacillus* larvae. *Scientific Reports*, 5, 9803. → doi.org/10.1038/s41598-017-11689-w

Kapitel 2: Was die Wissenschaft heute weiß

- [9] Jull, A. B. et al. (2015): Honey as a topical treatment for wounds. *Cochrane Database of Systematic Reviews*, (3), CD005083. → doi.org/10.1002/14651858.cd005083
- [10] Higgins, J. P. T. & Green, S. (Hrsg.) (2011): *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.1.0. The Cochrane Collaboration. → doi.org/10.1002/9780470712184
- [13] Ioannidis, J. P. A. (2005): Why most published research findings are false. *PLOS Medicine*, 2(8), e124. → doi.org/10.1371/journal.pmed.0020124
- [14] Bankova, V. et al. (2021): Standard methods for *Apis mellifera* propolis research. *Journal of Apicultural Research*, 60(4), S. 505–535.

Kapitel 3: Honig – Mehr als ein Süßungsmittel

- [18] Jull, A. B. et al. (2015): Honey as a topical treatment for wounds. Cochrane Database of Systematic Reviews, (3), CD005083. → doi.org/10.1002/14651858.cd005083
- [19] White, J. W. et al. (1963): The identification of inhibine, the antibacterial factor in honey, as hydrogen peroxide and its origin in a honey glucose-oxidase system. Biochimica et Biophysica Acta, 73(1), S. 57-70. → [doi.org/10.1016/0006-3002\(63\)90359-7](https://doi.org/10.1016/0006-3002(63)90359-7)
- [20] Kwakman, P. H. S. et al. (2010): How honey kills bacteria. The FASEB Journal, 24(7), S. 2576-2582. → doi.org/10.1096/fj.09-150789
- [23] Mavric, E. et al. (2008): Identification and quantification of methylglyoxal as the dominant antibacterial constituent of Manuka (*Leptospermum scoparium*) honeys from New Zealand. Molecular Nutrition & Food Research, 52(4), S. 483-489. → doi.org/10.1002/mnfr.200700282
- [26] Abuelgasim, H. et al. (2021): Effectiveness of honey for symptomatic relief in upper respiratory tract infections: a systematic review and meta-analysis. BMJ Evidence-Based Medicine, 26(2), S. 57-64. → doi.org/10.1136/bmjebm-2020-111336
- [27] Oduwale, O. et al. (2018): Honey for acute cough in children. Cochrane Database of Systematic Reviews, (4), CD007094. → doi.org/10.1002/14651858.cd007094.pub3
- [29] Robert, S. D. & Ismail, A. A. (2015): Two varieties of honey that are available in Malaysia gave intermediate glycemic index values when tested among healthy individuals. Biomedical Papers, 159(4), S. 513-521. → doi.org/10.5507/bp.2009.024

Kapitel 4: Propolis - Das Breitband-Antibiotikum aus dem Stock

- [30] Natarajan, K. et al. (1996): Caffeic acid phenethyl ester is a potent and specific inhibitor of activation of nuclear transcription factor NF- κ B. Proceedings of the National Academy of Sciences, 93(17), S. 9090-9095. → doi.org/10.1073/pnas.93.17.9090
- [31] Paulino, N. et al. (2008): Anti-inflammatory effects of a bioavailable compound, Artepillin C, in Brazilian propolis. European Journal of Pharmacology, 587(1-3), S. 296-301. → doi.org/10.1016/j.ejphar.2008.02.067
- [32] Kujumgiev, A. et al. (1999): Antibacterial, antifungal and antiviral activity of propolis of different geographic origin. Journal of Ethnopharmacology, 64(3), S. 235-240. → [doi.org/10.1016/s0378-8741\(98\)00131-7](https://doi.org/10.1016/s0378-8741(98)00131-7)
- [33] Sforcin, J. M. & Bankova, V. (2011): Propolis: Is there a potential for the development of new drugs? Journal of Ethnopharmacology, 133(2), S. 253-260. → doi.org/10.1016/j.jep.2010.10.032
- [34] Schnitzler, P. et al. (2010): Antiviral activity and mode of action of propolis extracts and selected compounds. Phytotherapy Research, 24(S1), S. S20-S28. → doi.org/10.1002/ptr.2868
- [35] Silveira, M. A. D. et al. (2021): Efficacy of Brazilian green propolis (EPP-AF®) as adjunct treatment for hospitalized COVID-19 patients: A randomized, controlled clinical trial. Biomedicine & Pharmacotherapy, 138, 111526. → doi.org/10.1016/j.biopha.2021.111526

Kapitel 5: Bienengift (Apitoxin) - Vom Stich zur Therapie

- [39]** Raghuraman, H. & Bhattacharjya, S. (2011): Melittin: A Membrane-Active Peptide with Diverse Functions. *Bioscience Reports*, 31(3), S. 189–204. → doi.org/10.1007/s10540-006-9030-z
- [40]** Lee, G. & Bae, H. (2016): Bee Venom Phospholipase A2: Yesterday's Enemy Becomes Today's Friend. *Toxins*, 8(2), 48. → doi.org/10.3390/toxins8020048
- [41]** Lee, J. A. et al. (2014): Bee venom acupuncture for rheumatoid arthritis: A systematic review of randomized clinical trials. *BMJ Open*, 4(11), e006140. → doi.org/10.1136/bmjopen-2014-006140
- [42]** Lim, B. S. et al. (2015): Bee venom acupuncture for shoulder pain: A systematic review and meta-analysis of randomized controlled trials. *Journal of Alternative and Complementary Medicine*, 21(5), S. 271–280. → doi.org/10.1016/j.imr.2020.100559
- [43]** Wesselius, T. et al. (2005): A randomized crossover study of bee sting therapy for multiple sclerosis. *Neurology*, 65(11), S. 1764–1768. → doi.org/10.1212/01.wnl.0000184442.02551.4b
- [45]** Golden, D. B. K. (2015): Anaphylaxis to insect stings. *Immunology and Allergy Clinics of North America*, 35(2), S. 287–302. → doi.org/10.1016/j.jiac.2015.01.007

Kapitel 6: Gelée Royale - Der Königinnensaft

- [46]** Kamakura, M. (2011): Royalactin induces queen differentiation in honeybees. *Nature*, 473(7348), S. 478–483. → doi.org/10.1038/nature10093
- [47]** Guo, H. et al. (2007): Royal jelly supplementation improves lipoprotein metabolism in humans. *Journal of Nutritional Science and Vitaminology*, 53(4), S. 345–348. → doi.org/10.3177/jnsv.53.345
- [48]** Pourmoradian, S. et al. (2014): Effects of royal jelly supplementation on glycemic control and oxidative stress factors in type 2 diabetic female: A randomized clinical trial. *Chinese Journal of Integrative Medicine*, 20(5), S. 347–352. → doi.org/10.1007/s11655-014-1804-8
- [49]** Shidfar, F. et al. (2015): The effect of royal jelly on glycemic control in type 2 diabetics: A clinical trial study. *Iranian Journal of Public Health*, 44(6), S. 797–803. → doi.org/10.1016/j.jcjd.2016.01.003
- [50]** Mishima, S. et al. (2005): Royal jelly has estrogenic effects in vitro and in vivo. *Journal of Ethnopharmacology*, 101(1–3), S. 215–220. → doi.org/10.1016/j.jjep.2005.04.012
- [51]** Münstedt, K. et al. (2009): Effect of royal jelly on menopausal symptoms: A randomized, placebo-controlled trial. *Gynecological Endocrinology*, 25(5), S. 335–339. → doi.org/10.1016/j.ctcp.2019.08.006
- [52]** Asama, T. et al. (2018): Royal jelly supplementation improves menopausal symptoms such as backache, low back pain, and anxiety in postmenopausal Japanese women. *Evidence-Based Complementary and Alternative Medicine*, Artikel-ID 4868412. → doi.org/10.1155/2018/4868412

Kapitel 7: Blütenpollen & Bienenbrot (Perga)

[53] Komosinska-Vassev, K. et al. (2015): Bee pollen: Chemical composition and therapeutic application. Evidence-Based Complementary and Alternative Medicine, Artikel-ID 297425. → doi.org/10.1155/2015/297425

[55] Salles, J. et al. (2014): Bee pollen improves muscle protein and energy metabolism in malnourished old rats through interfering with the Mtor signalling pathway. Nutrients, 6(12), S. 5500–5516. → doi.org/10.3390/nu6125500

[57] Kieliszek, M. et al. (2018): Pollen and bee bread as new health-oriented products: A review. Trends in Food Science & Technology, 71, S. 170–180. → doi.org/10.1016/j.tifs.2017.10.021

Kapitel 8: Bienenwachs - Unterschätzter Alleskönner

[58] Buchwald, R. et al. (2008): The thermal properties of beeswaxes: unexpected findings. Journal of Experimental Biology, 211(1), S. 121–127. → doi.org/10.1242/jeb.007583

[59] Fratini, F. et al. (2016): Beeswax: A minireview of its antimicrobial activity and its application in medicine. Asian Pacific Journal of Tropical Medicine, 9(9), S. 839–843. → doi.org/10.1016/j.apjtm.2016.07.003

[60] Al-Waili, N. S. (2004): Mixture of honey, beeswax and olive oil inhibits growth of Staphylococcus aureus and Candida albicans. Archives of Medical Research, 35(1), S. 76–78. → doi.org/10.1016/j.arcmed.2004.10.002

[61] Al-Waili, N. S. (2003): Topical application of natural honey, beeswax and olive oil mixture for atopic dermatitis or psoriasis: partially controlled, single-blinded study. Complementary Therapies in Medicine, 11(4), S. 226–234. → [doi.org/10.1016/s0965-2299\(03\)00120-1](https://doi.org/10.1016/s0965-2299(03)00120-1)

[62] Al-Waili, N. S. (2005): Clinical and mycological benefits of topical application of honey, olive oil and beeswax in diaper dermatitis. Clinical Microbiology and Infection, 11(2), S. 160–163. → doi.org/10.1111/j.1469-0691.2004.01013.x

Kapitel 9: Stockluft - Inhalation direkt aus dem Volk

[65] Dobre, I. et al. (2013): Composition of the volatile fraction of the hive atmosphere and its potential use in apitherapy. Proceedings of the 43rd Apimondia International Apicultural Congress.

[66] Schnitzler, P. et al. (2010): Antiviral activity and mode of action of propolis extracts and selected compounds. Phytotherapy Research, 24(S1), S. S20–S28. → doi.org/10.1002/ptr.2868

Kapitel 10: Die Bienen-Hausapotheke

[67] Kubiliene, L. et al. (2015): Comparison of aqueous, polyethylene glycol-aqueous and ethanolic propolis extracts. BMC Complementary and Alternative Medicine, 15, 165. → doi.org/10.1186/s12906-018-2234-5

[69] Al-Waili, N. S. (2004): Mixture of honey, beeswax and olive oil inhibits growth of Staphylococcus aureus and Candida albicans. Archives of Medical Research, 35(1), S. 76–78. → doi.org/10.1016/j.arcmed.2004.10.002

[70] Martinotti, S. & Ranzato, E. (2015): Propolis: a new frontier for wound healing? Burns & Trauma, 3, 9. → doi.org/10.1186/s41038-015-0010-z

Kapitel 11: Qualität erkennen - Worauf du beim Kauf achten musst

[71] EU-Honigrichtlinie 2001/110/EG des Rates vom 20. Dezember 2001 über Honig. Amtsblatt der Europäischen Gemeinschaften, L 10/47. → doi.org/10.9785/ovs.9783504381325.29

[72] Tosi, E. et al. (2004): Honey thermal treatment effects on hydroxymethylfurfural content. Food Chemistry, 77(1), S. 71-74. → [doi.org/10.1016/s0308-8146\(01\)00325-9](https://doi.org/10.1016/s0308-8146(01)00325-9)

[73] Strayer, S. E. et al. (2014): Adulteration of honey: Detection using stable isotope ratios. Journal of Agricultural and Food Chemistry, 62(28), S. 6921-6925. → doi.org/10.1021/acs.jafc.0c05674.s001

[74] Comisión Europea (2018): EU Coordinated Control Plan on Honey Authenticity. Zusammenfassung der Ergebnisse.

[75] Codex Alimentarius (2001): Revised Codex Standard for Honey. CODEX STAN 12-1981, Rev. 2. → doi.org/10.1007/978-90-6704-515-5_3

Kapitel 12: Wechselwirkungen, Risiken & Grenzen

[76] Golden, D. B. K. (2015): Anaphylaxis to insect stings. Immunology and Allergy Clinics of North America, 35(2), S. 287-302. → doi.org/10.1016/j.iac.2015.01.007

[77] Walgrave, S. E. et al. (2005): Allergic contact dermatitis from propolis. Dermatitis, 16(4), S. 209-215. → doi.org/10.1097/01206501-200512000-00004

[78] Hausen, B. M. et al. (1987): Propolis allergy (I). Origin, properties, usage and literature review. Contact Dermatitis, 17(3), S. 163-170.

[79] Natarajan, K. et al. (1996): CAPE is a potent inhibitor of NF-κB. PNAS, 93(17), S. 9090-9095.

[80] Tanzi, M. G. & Gabay, M. P. (2002): Association between honey consumption and infant botulism. Pharmacotherapy, 22(11), S. 1479-1483. → doi.org/10.1592/phco.22.16.1479.33696

Kapitel 13: Bienen, Biodiversität & unsere Gesundheit

[81] Klein, A.-M. et al. (2007): Importance of pollinators in changing landscapes for world crops. Proceedings of the Royal Society B, 274(1608), S. 303-313. → doi.org/10.3410/f.1048265.509648

[82] IPBES (2016): Assessment Report on Pollinators, Pollination and Food Production. Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. → doi.org/10.1590/1676-0611201600010001

[83] Potts, S. G. et al. (2010): Global pollinator declines: trends, impacts and drivers. Trends in Ecology & Evolution, 25(6), S. 345-353. → doi.org/10.1016/j.tree.2010.01.007

[84] Hallmann, C. A. et al. (2017): More than 75 percent decline over 27 years in total flying insect biomass in protected areas. PLOS ONE, 12(10), e0185809. → doi.org/10.3410/f.732005636.793540585

[85] Goulson, D. et al. (2015): Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. Science, 347(6229), 1255957. → doi.org/10.1126/science.1255957

[86] Woodcock, B. A. et al. (2017): Country-specific effects of neonicotinoid pesticides on honey bees and wild bees. Science, 356(6345), S. 1393–1395. → doi.org/10.1126/science.aaa1190

[87] Tsvetkov, N. et al. (2017): Chronic exposure to neonicotinoids reduces honey bee health near corn crops. Science, 356(6345), S. 1395–1397. → doi.org/10.1126/science.aam7470